

# CLASS TEST

S.No. : 05 BS\_CS\_A\_110919

Engineering Mathematics



## MADE EASY

India's Best Institute for IES, GATE & PSUs

Delhi | Noida | Bhopal | Hyderabad | Jaipur | Lucknow | Indore | Pune | Bhubaneswar | Kolkata | Patna

Web: [www.madeeasy.in](http://www.madeeasy.in) | E-mail: [info@madeeasy.in](mailto:info@madeeasy.in) | Ph: 011-45124612

# CLASS TEST 2019-2020

## COMPUTER SCIENCE & IT

Date of Test : 11/09/2019

### ANSWER KEY > Engineering Mathematics

1. (b)	7. (b)	13. (a)	19. (d)	25. (c)
2. (c)	8. (b)	14. (d)	20. (a)	26. (c)
3. (b)	9. (d)	15. (a)	21. (c)	27. (b)
4. (b)	10. (b)	16. (b)	22. (b)	28. (c)
5. (a)	11. (a)	17. (b)	23. (d)	29. (b)
6. (d)	12. (c)	18. (d)	24. (d)	30. (a)


## DETAILED EXPLANATIONS

1. (b)

$$\begin{aligned} f'(x) &= 6x^2 - 30x + 36 \\ &= 6(x-2)(x-3) \end{aligned}$$

So,  $f'(x) > 0$  when  $x < 2$  and also when  $x > 3$ .  $f(x)$  is increasing in  $]-\infty, 2[ \cup ]3, \infty[$ .

OR, by Wavy-Curve Method



2. (c)

$$\begin{aligned} A + A' &= \begin{bmatrix} \cos\alpha & \sin\alpha \\ -\sin\alpha & \cos\alpha \end{bmatrix} + \begin{bmatrix} \cos\alpha & -\sin\alpha \\ \sin\alpha & \cos\alpha \end{bmatrix} \\ &= \begin{bmatrix} 2\cos\alpha & 0 \\ 0 & 2\cos\alpha \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \end{aligned}$$

$$\therefore 2\cos\alpha = 1$$

$$\Rightarrow \alpha = \frac{\pi}{3}$$

3. (b)

$A$  is skew-symmetric,

$$\Rightarrow A = -A^T$$

$$\text{Now, } (A \cdot A)^T = A^T \cdot A^T = (-A) \cdot (-A) = A \cdot A$$

$\therefore A \cdot A$  is a symmetric matrix.

4. (b)

$$\lambda = np = \frac{1}{100} \times 100 = 1$$

$$P(X > 2) = 1 - (P(X=0) + P(X=1))$$

$$P(X=0) = \frac{e^{-\lambda} \cdot \lambda^0}{0!} = e^{-\lambda}$$

$$P(X=1) = \frac{e^{-\lambda} \lambda^1}{1!} = e^{-\lambda} \cdot \lambda$$

$$P(X > 2) = 1 - e^{-1}(2) = \frac{1-2}{e} = \frac{e-2}{e}$$

5. (a)

Given

$$P = \int_0^1 x e^x dx$$

$$= \left[ x \int e^x dx \right]_0^1 - \int_0^1 \left[ \frac{d}{dx}(x) \int e^x dx \right] dx$$

$$\begin{aligned}
 &= \left[ x e^x \right]_0^1 - \int_0^1 (1) e^x dx \\
 &= (e^1 - 0) - \left[ e^x \right]_0^1 \\
 &= e^1 - [e^1 - e^0] \\
 &= e - e + 1 = 1
 \end{aligned}$$

6. (d)

$$\begin{aligned}
 I &= \int_0^{\pi/4} \log\left(\frac{\sin x}{\cos x}\right) dx = \int_0^{\pi/4} [\log(\sin x) dx - \log(\cos x) dx] \\
 &= \int_0^{\pi/2} \log \sin\left(\frac{\pi}{2} - x\right) dx - \int_0^{\pi/2} \log(\cos x) dx \\
 I &= 0
 \end{aligned}$$

7. (b)

$$\begin{aligned}
 |A - \lambda I| &= 0 \\
 \left| \begin{bmatrix} 1 - \lambda & \sin x \\ \sin x & 1 - \lambda \end{bmatrix} \right| &= 0 \\
 (1 - \lambda)^2 - \sin^2 x &= 0 \\
 1 + \lambda^2 - 2\lambda - \sin^2 x &= 0 \\
 \lambda^2 - 2\lambda + \cos^2 x &= 0 \\
 \lambda &= \frac{2 \pm \sqrt{4 - 4\cos^2 x}}{2} \\
 \lambda &= 1 \pm \sin x
 \end{aligned}$$

8. (b)

$$\begin{aligned}
 \frac{\partial f}{\partial x} &= 2 - 2x & \frac{\partial f}{\partial y} &= 2 - 2y \\
 r = \frac{\partial^2 f}{\partial x^2} &= -2 & t = \frac{\partial^2 f}{\partial y^2} &= -2, & s = \frac{\partial^2 f}{\partial x \partial y} &= 0
 \end{aligned}$$

finding stationary points,

$$\frac{\partial f}{\partial x} = 2 - 2x = 0$$

$$\Rightarrow x = 1$$

$$\frac{\partial f}{\partial y} = 2 - 2y = 0$$

$$\Rightarrow y = 1$$

at the stationary point (1, 1)

$$rt - s^2 = (-2)(-2) - 0 = 4 > 0$$

So,  $f(x, y)$  is maxima at (1, 1)

$$\begin{aligned}
 \text{Maximum value of } f(x, y) &= 2 + 2 + 2 - 1 - 1 \\
 &= 4
 \end{aligned}$$

9. (d)

The constant term in any characteristic polynomial is always  $|A|$ .

So,  $|A| = -\frac{1}{4}$  since constant term of  $p(\lambda)$  is  $-\frac{1}{4}$ .

10. (b)

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} (x - 1) = 0$$

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (x^3 - 1) = 0$$

Also  $f(1) = 0$

Thus  $\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^+} f(x) = f(1)$

$\Rightarrow f$  is continuous at  $x = 1$

And  $Lf'(1) = 2, Rf'(1) = 1$

$\Rightarrow f$  is not differentiable at  $x = 1$

11. (a)

Total possible outcomes =  ${}^{52}C_2 = 1326$

Favourable outcomes = Drawing any spade apart from king of spades along with any king left in pack + Drawing king of spades with any three kings left in pack

**Note:** It is necessary that spade and king's card should be different. So in 2<sup>nd</sup> case, when king of spade's is drawn it is considered as a spade.

$\therefore$  Favourable outcomes =  ${}^{12}C_1 \times {}^4C_1 + {}^1C_1 \times {}^3C_1 = 51$

$$\text{Probability} = \frac{51}{1326} = \frac{1}{26}$$

12. (c)

$$\lim_{x \rightarrow \infty} \left( \frac{x}{2+x} \right)^{2x} = \lim_{x \rightarrow \infty} \left( \frac{2+x}{x} \right)^{-2x}$$

$$\lim_{x \rightarrow \infty} \left( 1 + \frac{2}{x} \right)^{-2x} = \lim_{x \rightarrow \infty} \left( 1 + \frac{2}{x} \right)^{\frac{x}{2}(-4)} \quad \because 2x = \frac{x}{2}(-4)$$

$$= e^{-4} \left( \because \lim_{x \rightarrow \infty} \left( 1 + \frac{1}{x} \right)^x = e \right)$$

13. (a)

$$a_{ij} = \begin{cases} i; & i = j \\ 0; & i \neq j \end{cases}$$

$$\Rightarrow [A] = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 2 & 0 & \dots & 0 \\ 0 & 0 & 3 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & n \end{bmatrix}$$

Determinant of all n eigen value of A

$$= \text{Product of diagonal elements}$$

$$= 1 \times 2 \times \dots \times n = n!$$

15. (a)

$[A : B]$

$$\left[ \begin{array}{cccc|c} 1 & 2 & -2 & 0 & 0 \\ 2 & -1 & -1 & 0 & 0 \\ 1 & 2 & -1 & 0 & 0 \\ 4 & -1 & -1 & 3 & 0 \end{array} \right]$$

$$R_4 \rightarrow R_4 - R_2, \quad R_3 \rightarrow (R_3 - R_1)$$

$$= \left[ \begin{array}{cccc|c} 1 & 2 & -2 & 0 & 0 \\ 2 & -1 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 2 & 0 & 0 & 3 & 0 \end{array} \right]$$

$$R_1 \rightarrow \frac{1}{3}(R_1 + 2R_2 + 4R_3)$$

$$= \left[ \begin{array}{cccc|c} 1 & 0 & 0 & 0 & 0 \\ 2 & -1 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 2 & 0 & 0 & 3 & 0 \end{array} \right]$$

$$R_2 \rightarrow (2R_1 - R_2 - R_3), \quad R_4 \rightarrow \frac{1}{3}(R_4 - 2R_1)$$


$$= \left[ \begin{array}{cccc|c} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{array} \right]$$

$$\rho(A : B) = \rho(A) = 4 = \text{number of variables}$$

$\Rightarrow$  System is consistent with trivial solution.

16. (b)

The tree diagram for problem is



$$P(R) = \frac{m}{m+n} \times \frac{n+1}{n+m+1} + \frac{n}{m+n} \times \frac{n}{n+m+1}$$

$$= \frac{m(n+1) + n^2}{(m+n)(n+m+1)}$$

17. (b)

$$\begin{aligned}
 \lim_{x \rightarrow 0} \frac{e^{ax} - e^{-ax}}{\log(1+bx)} &= \lim_{x \rightarrow 0} \frac{(e^{ax} - e^{-ax}) \times 2ax \times b}{2ax \times b \times \log(1+bx)} \\
 &= \lim_{x \rightarrow 0} \left( \frac{e^{ax} - e^{-ax}}{2ax} \right) \times \lim_{x \rightarrow 0} \frac{bx}{\log(1+bx)} \left( \frac{2a}{b} \right) \\
 &= \lim_{x \rightarrow 0} \left( \frac{\sinh ax}{ax} \right) \lim_{x \rightarrow 0} \frac{bx}{\log(1+bx)} \left( \frac{2a}{b} \right) \\
 &= 1 \times 1 \times \frac{2a}{b} \\
 &= \frac{2a}{b}
 \end{aligned}$$

18. (d)

$$\begin{aligned}
 y &= -\int \frac{1 - \sin x - 1}{1 - \sin x} dx \\
 &= -\int 1 \cdot dx + \int \frac{1}{1 - \sin x} dx \\
 y &= -x + \int \frac{dx}{1 - \sin x} \\
 \int \frac{dx}{1 - \sin x} &= \int \frac{1 + \sin x dx}{(1 - \sin x)(1 + \sin x)} = \int \frac{(1 + \sin x)}{(1 - \sin^2 x)} dx \\
 &= \int \frac{1 + \sin x}{\cos^2 x} dx \\
 &= \int \sec^2 x dx + \int \sec x \tan x dx \\
 &= \tan x + \sec x + C \\
 y &= -x + \tan x + \sec x + C
 \end{aligned}$$

19. (d)

$$AX = B$$

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 5 \\ 2 & 3 & \lambda \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 6 \\ 10 \\ \mu \end{bmatrix}$$

$$C = (A, B) = \begin{bmatrix} 1 & 1 & 1 & : & 6 \\ 1 & 2 & 5 & : & 10 \\ 2 & 3 & \lambda & : & \mu \end{bmatrix}$$

After performing  $R_2 \leftarrow R_2 - R_1$  and  $R_3 \leftarrow R_3 - 2R_1$

$$C = \begin{bmatrix} 1 & 1 & 1 & : & 6 \\ 0 & 1 & 4 & : & 4 \\ 0 & 1 & \lambda - 2 & : & \mu - 12 \end{bmatrix}$$

After performing  $R_3 \leftarrow R_3 - R_2$

$$C = \begin{bmatrix} 1 & 1 & 1 & : & 6 \\ 0 & 1 & 4 & : & 4 \\ 0 & 0 & \lambda - 6 & : & \mu - 16 \end{bmatrix}$$

Since  $R(A) = R(C)$  for unique solution

So  $\lambda - 6 \neq 0$ ,  $\lambda \neq 6$  and  $\mu - 10 \neq 0$ ,  $\mu \neq 16$ .

For no solution  $R(A) \neq R(C)$  then  $R(A) = 2$  and  $R(C) = 3$

$$\lambda - 6 = 0$$

$\Rightarrow \lambda = 6$  and  $\mu - 16 \neq 0 \Rightarrow \mu \neq 16$

For infinite solution  $R(A) = R(C) = 2$

then  $\lambda - 6 = 0$  and  $\mu - 16 = 0$

$$\lambda = 6 \text{ and } \mu = 16$$

So all of options are true.

20. (a)

$$|x-2| = \begin{cases} -(x-2); & x < 2 \\ (x-2); & x > 2 \end{cases}$$

$$\begin{aligned} \int_1^3 \frac{|x-2|}{x} dx &= \int_1^2 \frac{-(x-2)}{x} dx + \int_2^3 \frac{x-2}{x} dx \\ &= \int_1^2 \left(-1 + \frac{2}{x}\right) dx + \int_2^3 \left(1 - \frac{2}{x}\right) dx \\ &= -(2-1) + (2\ln x)_1^2 + (x)_2^3 - 2(\ln x)_1^3 \\ &= 2\ln 2 - 2\ln \frac{3}{2} \\ &= 2\ln \frac{2}{3} = 2\ln \frac{4}{3} \\ &= 0.575 \end{aligned}$$

21. (c)

The system may be written in matrix form as

$$\begin{bmatrix} 1 & 3 & -8 \\ 1 & 4 & 3 \\ 1 & 3 & 4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 4 \\ -2 \\ 1 \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & 3 & -8 \\ 1 & 4 & 3 \\ 1 & 3 & 4 \end{bmatrix}$$

$$LU = A$$

$$= \begin{bmatrix} 1 & 3 & -8 \\ 1 & 4 & 3 \\ 1 & 3 & 4 \end{bmatrix}$$

$$\begin{aligned}
 l_{11} &= 1, l_{21} = l, l_{31} = l \\
 l_{11} u_{12} &= 3 \Rightarrow u_{12} = 3, \\
 l_{21} u_{12} + l_{22} &= 4 \Rightarrow l_{22} = 4 - 1 \cdot 3 = 1 \\
 l_{31} u_{12} + l_{32} &= 3 \Rightarrow l_{32} = 3 - 1 \times 3 = 0
 \end{aligned}$$

$$l_{11} u_{13} = -8 \Rightarrow u_{13} = \frac{-8}{1} = -8$$

$$\begin{aligned}
 l_{21} u_{13} + l_{22} u_{23} &= 3 \Rightarrow u_{23} = 11 \\
 l_{31} u_{13} + l_{32} u_{23} + l_{33} &= 4 \Rightarrow l_{33} = 12
 \end{aligned}$$

$$\therefore L = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 0 & 12 \end{bmatrix} \text{ and } U = \begin{bmatrix} 1 & 3 & -8 \\ 0 & 1 & 11 \\ 0 & 0 & 1 \end{bmatrix}$$

22. (b)

$$\begin{aligned}
 \lambda_1^2 + \lambda_2^2 + \lambda_1 \lambda_2 &= \lambda_1^2 + \lambda_2^2 + 2\lambda_1 \lambda_2 - \lambda_1 \lambda_2 \\
 &= (\lambda_1 + \lambda_2)^2 - \lambda_1 \lambda_2
 \end{aligned}$$

$$\begin{aligned}
 \text{Sum of eigen values, } \lambda_1 + \lambda_2 &= \text{trace of matrix} \\
 &= \text{sum of diagonal elements} \\
 &= 1 - \frac{1}{3} = \frac{2}{3}
 \end{aligned}$$

$$\begin{aligned}
 \text{Products of eigen values, } \lambda_1 \lambda_2 &= \text{determinant of matrix} \\
 &= 1 \left( -\frac{1}{3} \right) - (-1) \left( \frac{4}{9} \right) \\
 &= \frac{1}{9}
 \end{aligned}$$

$$\begin{aligned}
 \therefore (\lambda_1 + \lambda_2)^2 - \lambda_1 \lambda_2 &= \left( \frac{2}{3} \right)^2 - \frac{1}{9} \\
 &= \frac{4}{9} - \frac{1}{9} = \frac{3}{9} = 0.33
 \end{aligned}$$

23. (d)

$$\begin{aligned}
 6(13 \times 11 - 4 \times 37) - 3(32 \times 11 - 10 \times 37) + 7(32 \times 4 - 10 \times 13) \\
 = -30 + 54 - 14 \\
 = 10
 \end{aligned}$$

24. (d)

$$|A - \lambda I| = \begin{vmatrix} 1-\lambda & 1 & 3 \\ 1 & 5-\lambda & 1 \\ 3 & 1 & 1-\lambda \end{vmatrix} = 0$$

$$\lambda^3 - 7\lambda^2 + 36 = 0$$

since  $\lambda = 3$  is root of the equation

$$(\lambda - 3)(\lambda^2 - 4\lambda - 12) = 0$$

$$(\lambda - 3)(\lambda + 2)(\lambda - 6) = 0$$



highest eigen value = 6

$$(A - \lambda I)X = 0$$

for  $\lambda = 6$

$$\begin{bmatrix} -5 & 1 & 3 \\ 1 & -1 & 1 \\ 3 & 1 & -5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = 0$$

$$-5x_1 + x_2 + 3x_3 = 0, \quad x_1 - x_2 + x_3 = 0, \quad 3x_1 + x_2 - 5x_3 = 0$$

$$\frac{x_1}{4} = \frac{x_2}{8} = \frac{x_3}{4} \quad \text{or} \quad \frac{x_1}{1} = \frac{x_2}{2} = \frac{x_3}{1}$$

so eigen vector is  $[1, 2, 1]^T$

25. (c)

Using Crout's method

$$A = \begin{bmatrix} l_{11} & 0 \\ l_{21} & l_{22} \end{bmatrix} \begin{bmatrix} 1 & u_{12} \\ 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 4 \\ 6 & 3 \end{bmatrix} = \begin{bmatrix} l_{11} & 0 \\ l_{21} & l_{22} \end{bmatrix} \begin{bmatrix} 1 & u_{12} \\ 0 & 1 \end{bmatrix}$$

$$l_{11} = 2$$

$$l_{11}u_{12} = 4$$

$$u_{12} = \frac{4}{2} = 2$$

$$l_{21} = 6$$

$$l_{21}u_{12} + l_{22} = 3$$

$$6 \times 2 + l_{22} = 3$$

$$l_{22} = 3 - 12$$

$$l_{22} = -9$$

So, LU decomposition of given matrix is

$$L = \begin{bmatrix} 2 & 0 \\ 6 & -9 \end{bmatrix} \quad U = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$$

26. (c)

$$A^2 = \begin{bmatrix} \alpha & \beta \\ \beta & \alpha \end{bmatrix} = \begin{bmatrix} a & b \\ b & a \end{bmatrix} \begin{bmatrix} a & b \\ b & a \end{bmatrix}$$

$$= \begin{bmatrix} a^2 + b^2 & 2ab \\ 2ab & a^2 + b^2 \end{bmatrix}$$

$$\alpha = a^2 + b^2, \beta = 2ab$$

27. (b)

Consider  $n = 3$ 

Then 
$$A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ 3 & 6 & 9 \end{bmatrix}$$

and 
$$|A| = \begin{vmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ 3 & 6 & 9 \end{vmatrix} \begin{array}{l} R_3 \leftarrow 3R_1 - R_3 \\ R_2 \leftarrow 2R_1 - R_2 \end{array}$$

$$= \begin{vmatrix} 1 & 2 & 3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{vmatrix}$$

This if  $n = 3$  then Rank  $(A) = 1$ .

28. (c)

Given function, 
$$f(x) = 2x^3 - 15x^2 + 36x + 1$$
  

$$f'(x) = 6x^2 - 30x + 36$$
  

$$= 6(x-2)(x-3)$$

Hence,  $f'(x) > 0$  when  $x < 2$  and  $x > 3$ . So  $f(x)$  is increasing in  $]-\infty, 2[ \cup ]3, \infty[$ 

29. (b)

$$\begin{aligned} |\text{adj}(\text{adj } A)| &= |A|^{(n-1)^2} \\ |\text{adj}(\text{adj } A^2)| &= |A^2|^{(n-1)^2} \\ &= |A^2|^{(3-1)^2} = |A|^{2 \times (4)} \\ &= |A|^8 \end{aligned}$$

30. (a)

Let 
$$A = \lim_{x \rightarrow \pi/2} \frac{\cos x}{\left(x - \frac{\pi}{2}\right)^3}$$

By putting  $\left(x - \frac{\pi}{2}\right) = t$

when  $x \rightarrow \frac{\pi}{2}$ ,  $t \rightarrow 0$ 

then, 
$$A = \lim_{t \rightarrow 0} \frac{\cos\left(\frac{\pi}{2} + t\right)}{t^3}$$
  

$$= \lim_{t \rightarrow 0} \frac{-\sin t}{t^3} = \lim_{t \rightarrow 0} (-1) \frac{\sin t}{t} \cdot \frac{1}{t^2}$$
  

$$= (-1) \cdot 1 \cdot \frac{1}{0} = -\infty$$

